

IGSF8 is an Innate Immune Checkpoint and Cancer Immunotherapy Target


Karim Benhadji, MD

Cambridge, USA 21 October 2023



Resistance to Immune-Checkpoints and innate immunity


- Over 65% of tumors have antigen presentation defects and these tumors are often immune cold and don't respond to immunotherapy
- Innate immune Natural Killer (NK) cells normally kill cells with antigen presentation defects
- Why aren't tumors with MHC-I defects killed by NK cells?






IGSF8 was discovered through antibody Al ranking and CRISPR screens

CRISPR screen in cancer cells grown in syngeneic mouse tumors




CRISPR screens of cancer cells cocultured with primary NK cells






IGSF8 is overexpressed in solid tumors (TCGA)






IGSF8 is amplified in various solid tumors (TCGA)







IGSF8 mRNA expression is correlated with cold tumor markers





High IGSF8 mRNA is associated with poor survival and non-response to ICI






IGSF8 has receptors on both NK cells and Dendritic Cells (DCs) in human and mouse









Anti-IGSF8 leads to potent tumor growth inhibition across multiple tumor models as single agent and in combination





Anti-IGSF8 prolongs survival in B16 model and increases T, NK and dendritic cells infiltration








Conclusions

- IGSF8 is highly expressed on cancer cells
- Anti-IGSF8 enhances NK killing, antigen presentation, and turns immune-cold tumors hot
- Anti-IGSF8 shows monotherapy efficacy and is synergistic with anti-PD1 in multiple syngeneic tumor models (including some known to resist to anti-PD1)
- The IGSF8 inhibitor, GV20-0251 is currently explored in a phase 1 study in patients with advanced or metastatic solid tumors (NCT05669430)









Via Ginevra 4, CH-6900 Lugano T. +41 (0)91 973 19 00 esmo@esmo.org

esmo.org

